



IMHOTEP

AFRICAN JOURNAL OF PURE AND APPLIED MATHEMATICS

Imhotep Mathematical Proceedings Volume 1, Numéro 1, (2014), pp. 59 – 63.

Construction of Some New Three Associate Class Partially Balanced
Incomplete Block Designs in Two Replicates

E. C. Kipkemoi
kipched@gmail.com

J. k. Koske
koske4@co.uk

J. M. Mutiso
johnkasome@yahoo.com


Department of Statistics and
Computing, Moi University, P O
Box 3900, Eldoret, Kenya.

Abstract

Search for experimental designs which aid in research studies involving large number of treatments with minimal experimental units has been desired overtime. This paper constructs some new series of three associate Partially Balanced Incomplete Block (PBIB) designs having $n(n-2)/4$ treatments with three associate classes in two replicates using the concept of triangular association scheme. The design is constructed from an even squared array of n rows and n columns ($n \geq 8$) with its both diagonal entries bearing no treatment entries and that given the location of any treatment in the squared array, the other location of the same treatment in the array is predetermined. The design and association parameters for a general case of an even integer $n \geq 8$ are obtained with an illustrated case for $n = 8$. Efficiencies of the designs within the class of designs are obtained for a general case of even $n \geq 8$ with a listing of efficiencies of designs with blocks sizes in the interval $[8, 22]$. The designs constructed have three associate classes and are irreducible to minimum number of associate classes.

Proceedings of the 2nd Strathmore
International Mathematics Conference
(SIMC 2013), 12 - 16 August 2013,
Strathmore University, Nairobi, Kenya.

<http://imhotep-journal.org/index.php/imhotep/>

Imhotep Mathematical Proceedings 

Construction of Some New Three Associate Class Partially Balanced Incomplete Block Designs in Two Replicates

E. C. Kipkemoi, J. k. Koske and J. M. Mutiso

Abstract. Search for experimental designs which aid in research studies involving large number of treatments with minimal experimental units has been desired overtime. This paper constructs some new series of three associate Partially Balanced Incomplete Block (PBIB) designs having $n(n-2)/4$ treatments with three associate classes in two replicates using the concept of triangular association scheme. The design is constructed from an even squared array of n rows and n columns ($n \geq 8$) with its both diagonal entries bearing no treatment entries and that given the location of any treatment in the squared array, the other location of the same treatment in the array is predetermined. The design and association parameters for a general case of an even integer $n \geq 8$ are obtained with an illustrated case for $n = 8$. Efficiencies of the designs within the class of designs are obtained for a general case of even $n \geq 8$ with a listing of efficiencies of designs with blocks sizes in the interval [8,22]. The designs constructed have three associate classes and are irreducible to minimum number of associate classes.

Keywords. Partially Balanced Incomplete Block (PBIB), Associate class, three associate classes.

Introduction

By changing the arrangement of treatments or omitting certain blocks and or treatments, we obtain designs that may belong to a class of new designs. Using this technique Bose and Nair (1939) introduced some PBIB designs. Atiqullah (1958) established that considering PBIB designs based on triangular association scheme with $v = n((n-2))/2$, $b = (n-1)((n-2))/2$, $r = k = n-2$, $\lambda_1 = 1$ and $\lambda_2 = 2$. Other methods were given by Shrikande (1960, 1965) and Chang et al (1965) based on the existence of certain BIB designs and considering the dual of a BIB design as omitting certain blocks from the BIB design. John (1966) showed that triangular association scheme can be described by representing the treatments by ordered pairs (x,y) with $1 \leq x < y \leq q$ and then generalized (John, 1966) for the case that $v = q(q-1)((q-2))/6$, ($q > 3$). Arya and Narain (1981) discussed a new association scheme called truncated triangular (TT) with five associate classes when $v = p((p-2))/2$ with p an even positive integer ≥ 8 , and used to construct partial diallel crosses. Ching-Shui et al (1984) came up with a general and simple method of construction based on the relation of triangular and 1_2 type of PBIB design

Paper presented at the 2nd Strathmore International Mathematics Conference (SIMC 2013), 12 - 16 August 2013, Strathmore University, Nairobi, Kenya.

to the line graph theory. A construction of triangular designs with nested rows and columns is given by Agrawal and Prasad (1984). Sinha and Sanpei (2004) gave a series of triangular (tenary) designs with nested row and columns. Garg et al (May, 2011) constructed some new series of triangular and four associate PBIB designs with two replications by using dualization technique. Silver et al (April 2012) constructed some new three associate class Pbib Design with two replicates. Recently, Kipkemoi et al (January-June, 2013) constructed a three associate class partially balanced incomplete Block designs in two replicates

Construction of designs

The design is constructed from a squared array of n rows and n columns (n is even positive integer ≥ 8) with both diagonal entries in the array having no treatments allocated to as illustration in the figure below

*	n_{12}	n_{13}	n_{14}	n_{15}	n_{16}	n_{17}	*
n_{21}	*	n_{23}	n_{24}	n_{25}	n_{26}	*	n_{28}
n_{31}	n_{32}	*	n_{34}	n_{35}	*	n_{37}	n_{38}
n_{41}	n_{42}	n_{45}	*	*	n_{46}	n_{47}	n_{48}
n_{51}	n_{52}	n_{53}	*	*	n_{56}	n_{57}	n_{58}
n_{61}	n_{62}	*	n_{64}	n_{65}	*	n_{67}	n_{68}
n_{71}	*	n_{73}	n_{74}	n_{75}	n_{76}	*	n_{78}
*	n_{82}	n_{83}	n_{84}	n_{85}	n_{86}	n_{87}	*

i and j are integers ($1 \leq i, j \leq n$) such that;

Each row and column of the square array has $n - 2$ treatment entries.

The treatment entries are allocated in the array by following two subsequent steps

1. The initial set of v treatment entries are first filled on one triangle enclosed by the two diagonal and a side of the square array.
2. The second set of v treatment entries are replicated in each of the remaining triangles by simply reflecting the initial set of v treatment entries subsequently with the diagonal blank entries as mirrors in such a way that given any two entries n_{ij} and n_{ij} are allocated to treatment x if and only if the subscripts $i + i' = j + j'$ or $i + i' + j + j' = 2(n + 1)$.

Taking each row and column to constitute a block we obtain $n/2$ distinct blocks and thus a design with parameters

$$v = (n(n - 2))/4 \quad b = n/2 \quad k = n - 2 \quad r = 2 \quad \lambda_1 = 2 \quad \lambda_2 = 1 \quad \lambda_3 = 0$$

Association scheme

Two treatments are said to be:

- First associates if they both occur in the same row and column.
- Second associates if they both occur in the same row or the same column but not both.
- Third associates if they neither occur in the same row nor in the same column. Giving rise to the following association parameters.

$$n_1 = 1 \quad n_2 = 2(n - 4) \quad n_3 = \frac{n(n-10)+24}{4}$$

$$P_0 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 2(n-4) & 0 \\ 0 & 0 & 0 & \frac{n(n-10)+24}{4} \end{pmatrix} \quad P_1 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 2(n-4) & 0 \\ 0 & 0 & 0 & \frac{n(n-10)+24}{4} \end{pmatrix}$$

$$P_2 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 1 & 1 & n-4 & n-6 \\ 0 & 0 & n-6 & \frac{n(n-14)+48}{4} \end{pmatrix} \quad P_3 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 8 & 2(n-8) \\ 1 & 1 & 2(n-8) & \frac{n(n-18)+80}{4} \end{pmatrix}$$

Illustration: taking $n = 8$ we obtain three associate class PBIB design with the parameters

$$v = 12 \quad b = 4 \quad k = 6 \quad r = 2$$

$$\lambda_1 = 2 \quad \lambda_2 = 1 \quad \lambda_3 = 0$$

$$n_1 = 1 \quad n_2 = 8 \quad n_3 = 2$$

Whose blocks are:

1. (1, 2, 3, 4, 5, 6)
2. (1, 6, 7, 8, 9, 10)
3. (2, 5, 7, 10, 11, 12)
4. (3, 4, 8, 9, 11, 12)

With the parameters of the second kind given by:

$$P_0 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 8 & 0 \\ 0 & 0 & 0 & 2 \end{pmatrix} \quad P_1 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 8 & 0 \\ 0 & 0 & 0 & 2 \end{pmatrix}$$

$$P_2 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 8 & 0 \\ 0 & 0 & 0 & 2 \end{pmatrix} \quad P_3 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 8 & 0 \\ 1 & 1 & 0 & 0 \end{pmatrix}$$

The efficiency factors for this class of designs are given by

$$E_1 = \frac{n-4}{n-2}$$

$$E_2 = -\frac{n(n-4)^2}{7n^2-6n-16-n^3}$$

$$E_3 = -\frac{n(n-4)^2}{6n^2+2n-32-n^3}$$

and the overall efficiency factor of the design is

$$E = \frac{1}{n(n-2)-4} \left(\frac{4(n-4)}{n-2} - \frac{8n(n-4)^3}{7n^2-6n-16-n^3} + \frac{n(n-6)(n-4)^3}{6n^2+2n-32-n^3} \right)$$

Efficiencies of three associate class PBIB designs having $(n(n-2))/4$ treatments with two replicates for $6 \leq k \leq 22$ are give in the table below

n	v	b	k	r	λ_1	λ_2	λ_3	E	E₁	E₂	E₃
8	12	4	6	2	2	1	0	0.9495	0.6667	1	0.8889
10	20	5	8	2	2	1	0	0.9201	0.7500	0.9574	0.8738
12	30	6	10	2	2	1	0	0.9164	0.8000	0.9505	0.8807
14	42	7	12	2	2	1	0	0.9187	0.8333	0.9511	0.8906
16	56	8	14	2	2	1	0	0.9226	0.8571	0.9536	0.9000
18	72	9	16	2	2	1	0	0.9269	0.8750	0.9566	0.9083
20	90	10	18	2	2	1	0	0.9311	0.8889	0.9595	0.9156
22	110	11	20	2	2	1	0	0.9350	0.9000	0.9622	0.9219
24	132	12	22	2	2	1	0	0.9386	0.9091	0.9646	0.9274

Conclusion

In this paper we have constructed some new series of three associate class PBIB designs having $(n(n-2))/4$ treatments with two replications. The restriction of the number of replications to two helps to minimize cost. The average efficiency factors of these designs along with the three efficiencies factors E_1 , E_2 and E_3 are quite high for practical purposes.

References

- [1] H. Agrawal and J. Prasad, *Construction of partially balanced incomplete block designs with nested rows and columns*, *Biom. J.* **26**, (1984), 883 – 891.
- [2] A. S. Arya and P. Narain, *Truncated triangular association scheme and related partial diallel crosses*, *Sankhya: Indian journal of statistics* **43 B**, Pt 1, (1981), 93 – 103.
- [3] M. Atiqulla, *On configuration and non-isomorphism of some incomplete block designs*, *Indian journal of statistics*, 20 series 3, 4, (1958), 227 – 248.
- [4] R. C. Bose, K. R. Nair, *Partially Balanced Incomplete Block designs* *Sankhya* **4**, (1939), 337 – 372.
- [5] L. Chang, L. Changwen and L. Wan Ru, *incomplete block designs with triangular parameters for $k \geq 10$ and $v \geq 10$* , *Scientia sinica* **13**, (1965) 1493 – 1495
- [6] C. Ching-Shui, G.M. Constance and A.S. Hedayat *A unified method for constructing PBIB designs based on triangular and L2-schemes*, *J. R. Statist. Soc* **46**, 1,(1984), 31 – 37.
- [7] D. K. Garg, H.S. Jhaji and G. Mishra, *Construction of Some New Triangular and Four Associate Class PBIB Designs with Two Replicates*, *International Journal of Mathematical Sciences and Applications* **1**, **2**, (2011), 808 – 821.
- [8] S. Kishore and K. Sanpei, *Some series of block designs with nested rows and columns*, *Australasian Journal of combinatorics*, **29**, (2004), 337 – 347.
- [9] P. W. M. John, *An extension of the triangular association scheme to three associate classes*, *J. Roy. Statist. Soc B*, **28**, (1966), 361 – 365.
- [10] S. S. Shrikhande, *Relations between certain incomplete block designs*, *Contributions to Probability and Statistics*. I. Olkin (ed.). Stanford, CA: Stanford University Press, (1960), 388 – 395.
- [11] S. S. Shrikhande, *On a class of partially balanced incomplete block designs*, *Ann. Math. Statist.* **36**, (1965), 1807 – 1814.
- [12] J. K. R. Silver, E. C. Kipkemoi and I. K. Tum, *Construction of some new three associate class Pbib Designs with two replicates*, *International journal of academic research in progressive education and development* **1**, **2**, (2012), 188 – 192.
- [13] E. C. Kipkemoi, J. K. Koske and J. M. Mutiso, *Construction of three associate class partially balanced incomplete block designs in two replicates*, *American journal of Mathematical Science and Applications*, **1** (2013), 61 – 65.

E. C. Kipkemoi

e-mail: kipched@gmail.com

Department of Statistics and Computing, Moi University, P O Box 3900, Eldoret, Kenya.

J. k. Koske

e-mail: koske4@co.uk

Department of Statistics and Computing, Moi University, P O Box 3900, Eldoret, Kenya.

J. M. Mutiso

e-mail: johnkasome@yahoo.com

Department of Statistics and Computing, Moi University, P O Box 3900, Eldoret, Kenya.